

- COLLIN, S., DURANT, F. & EVRARD, G. (1986). *Acta Cryst.* **C42**, 199–201.
- COLLIN, S., EVRARD, G. & DURANT, F. (1986). *J. Cryst. Spectrosc. Res.* **16**, 255–269.
- DURANT, F., DE BEYS, V., COLLIN, S. & EVRARD, G. (1986). *Bull. Soc. Chim. Fr.* No. 4, pp. 497–503.
- JALFRE, M., BUCHER, B., DORME, N., MOCQUET, G. & PORSOLT, R. D. (1983). *Arch. Int. Pharmacodyn. Ther.* **264**, 232–256.
- LEJEUNE, J., MICHEL, A. & VERCAUTEREN, D. P. (1984). Program for interactive and flexible molecular fitting. Univ. of Namur, Belgium.
- MAIN, P., FISKE, S. J., HULL, S. E., LESSINGER, L., GERMAIN, G., DECLERCQ, J.-P. & WOOLFSON, M. M. (1980). *MULTAN80. A System of Computer Programs for the Automatic Solution of Crystal Structures from X-ray Diffraction Data*. Univs. of York, England, and Louvain, Belgium.
- SHELDRIK, G. M. (1976). *SHELX76*. Program for crystal structure determination. Univ. of Cambridge, England.
- STEWART, J. M., MACHIN, P. A., DICKINSON, C. W., AMMON, H. L., HECK, H. & FLACK, H. (1976). The *XRAY76* system. Tech. Rep. TR-446. Computer Science Center, Univ. of Maryland, College Park, Maryland.

Acta Cryst. (1987). **C43**, 577–579

Structure of (1R)-1,8,8-Trimethyl-3-oxabicyclo[3.2.1]octane-2,4-dione (Camphoric Anhydride)

BY KLAUS WICHMANN AND HANS BRADACZEK

Institute of Crystallography, Free University of Berlin, D-1000 Berlin 33, Federal Republic of Germany

AND ZBIGNIEW DAUTER AND TADEUSZ POŁOŃSKI

Institute of Organic and Food Chemistry and Technology, Technical University, 80-952 Gdańsk, Poland

(Received 12 June 1986; accepted 22 October 1986)

Abstract. $C_{10}H_{14}O_3$, $M_r = 182.22$, orthorhombic, $P2_12_12_1$, $a = 6.486$ (1), $b = 11.203$ (2), $c = 13.027$ (2) Å, $V = 946.6$ Å³, $Z = 4$, $D_x = 1.278$ (1) Mg m⁻³, $\lambda(\text{Cu } K\alpha) = 1.5418$ Å, $\mu(\text{Cu } K\alpha) = 0.778$ mm⁻¹, $F(000) = 392$, $T = 290$ (1) K, $R = 0.034$ for 905 observed independent reflections. The six-membered ring shows a half-boat conformation with marked non-planarity of the anhydride moiety.

Introduction. Knowledge of the ring geometry is indispensable for interpretation of the NMR spectra of glutaric anhydrides (Koer, de Hoog & Altona, 1975). The existence of large differences among ring torsion angles points to the strong influence of the substituents on the conformation of these compounds (Hašek, 1985). Changes of the ring and chromophore geometry are responsible for the strong substituent and solvent dependence of circular dichroism (CD) spectra of cyclic anhydrides. Interesting examples of this are the camphoric acid derivatives with a bicyclic skeleton for which the CD curves are strongly solvent dependent (Połoński, 1983). The structure of the bicyclic title compound was required for the analysis of its CD spectrum.

Experimental. Crystals of the title compound were grown from toluene. Unit-cell parameters were determined from oscillation and Weissenberg photographs, which showed orthorhombic symmetry with systematic

absences $h00$ for h odd, $0k0$ for k odd and $00l$ for l odd, indicating space group $P2_12_12_1$. A needle-like crystal of dimensions $0.75 \times 0.25 \times 0.20$ mm was used for data collection on a Stoe automatic four-circle diffractometer. Accurate cell parameters were obtained from a least-squares fit of θ values for 43 high-order reflections. The following data collection regime was applied: $\omega-2\theta$ scan, 55 steps of 0.2° , count time per step $0.3-1.8$ s, $\theta_{\max} = 60^\circ$ ($0 \leq h \leq 7$, $0 \leq k \leq 12$, $0 \leq l \leq 14$), two standard intensities (006 and 040) were checked every 90 min with no significant variation of their intensities. 926 unique reflections were measured of which 21 with $I < 2\sigma(I)$ were treated as unobserved in subsequent calculations. Lorentz and polarization but no absorption corrections were applied.

The structure was solved by direct methods and refined by full-matrix least squares using the *XTAL* package of programs (Stewart & Hall, 1983) on a VAX computer. The correct enantiomer was chosen on the basis of the known chirality of natural camphor, from which the compound was synthesized. H-atom positions were taken from a difference Fourier map and refined individually with isotropic temperature factors. Atomic scattering factors were taken from *International Tables for X-ray Crystallography* (1974). Minimized function $\sum w(F_o - F_c)^2$, with final values of $R = 0.034$ and $wR = 0.033$; $w = 1/\sigma^2(F)$. Ratio of maximum parameter shift to e.s.d. in the last cycle of refinement was 0.02 for non-hydrogen atoms and 0.05

for hydrogens. Maximum and minimum heights in final $\Delta\rho$ map were 0.15 and $-0.17 \text{ e} \text{ \AA}^{-3}$. The final atomic coordinates are listed in Table 1.*

Discussion. The bond lengths and angles are listed in Table 2. The molecular structure is shown with the atom numbering scheme in Fig. 1 and unit-cell contents are shown in Fig. 2.

All bond lengths agree well with standard values for corresponding bond types in glutaric anhydrides of known structure: bromovalerone anhydride (Hoehne, 1966); β -chloroglutaric anhydride (Koer *et al.*, 1975); *meso*-2,4-dimethylglutaric anhydride (Haltiwanger, Walba & Wand, 1980); *erythro*-2-methoxymethyl-2,4-dimethylglutaric anhydride (Hašek, 1985); α -phenylglutaric anhydride (Bocelli & Grenier-Loustalot, 1982)

* Lists of structure factors, anisotropic thermal parameters, H-atom coordinates, torsion angles and least-squares planes have been deposited with the British Library Document Supply Centre as Supplementary Publication No. SUP 43516 (13 pp.). Copies may be obtained through The Executive Secretary, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

Table 1. Final positional parameters ($\times 10^4$) and equivalent isotropic thermal parameters ($\times 10^2$) for non-hydrogen atoms with their *e.s.d.*'s in parentheses

U_{eq} is defined according to Hamilton (1959).

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{eq}} (\text{\AA}^2)$
C(1)	823 (5)	-281 (2)	252 (2)	4.58
O(1)	4102 (3)	784 (2)	553 (2)	5.34
C(2)	3050 (5)	-71 (3)	-12 (2)	5.12
O(2)	4010 (4)	-537 (2)	-688 (2)	8.24
C(3)	3325 (5)	1260 (3)	1456 (2)	5.05
O(3)	4462 (4)	1873 (2)	1942 (2)	7.52
C(4)	1128 (4)	980 (2)	1685 (2)	4.48
C(5)	-213 (5)	1692 (3)	918 (3)	5.81
C(6)	-351 (5)	889 (3)	-27 (2)	5.63
C(7)	667 (4)	-336 (2)	1436 (2)	4.24
C(8)	2183 (5)	-1206 (3)	1937 (3)	5.62
C(9)	-1532 (5)	-648 (4)	1793 (3)	6.08
C(10)	23 (7)	-1373 (4)	-326 (3)	7.38




Fig. 1. Perspective view of the molecule.

Table 2. Bond lengths (\AA) and angles ($^\circ$)

C(1)–C(2)	1.503 (4)	C(3)–O(3)	1.189 (4)
C(1)–C(6)	1.559 (4)	C(3)–C(4)	1.490 (4)
C(1)–C(7)	1.547 (3)	C(4)–C(5)	1.546 (4)
C(1)–C(10)	1.527 (5)	C(4)–C(7)	1.539 (4)
O(1)–C(2)	1.387 (4)	C(5)–C(6)	1.527 (5)
O(1)–C(3)	1.388 (4)	C(7)–C(8)	1.530 (4)
C(2)–O(2)	1.198 (4)	C(7)–C(9)	1.541 (4)
C(2)–C(1)–C(6)	106.5 (2)	O(3)–C(3)–C(4)	127.5 (3)
C(2)–C(1)–C(7)	107.3 (2)	C(3)–C(4)–C(5)	107.5 (2)
C(2)–C(1)–C(10)	109.9 (3)	C(3)–C(4)–C(7)	110.2 (2)
C(6)–C(1)–C(7)	103.5 (2)	C(5)–C(4)–C(7)	104.4 (2)
C(6)–C(1)–C(10)	113.1 (3)	C(4)–C(5)–C(6)	104.5 (3)
C(7)–C(1)–C(10)	115.9 (3)	C(1)–C(6)–C(5)	106.2 (3)
C(2)–O(1)–C(3)	122.5 (2)	C(1)–C(7)–C(4)	99.2 (2)
C(1)–C(2)–O(1)	117.4 (2)	C(1)–C(7)–C(8)	114.1 (2)
C(1)–C(2)–O(2)	126.8 (3)	C(1)–C(7)–C(9)	111.8 (3)
O(1)–C(2)–O(2)	115.8 (3)	C(4)–C(7)–C(8)	113.3 (2)
O(1)–C(3)–O(3)	116.6 (3)	C(4)–C(7)–C(9)	109.5 (3)
O(1)–C(3)–C(4)	115.8 (2)	C(8)–C(7)–C(9)	108.8 (3)




Fig. 2. View of the molecules within the unit cell.

and 1-oxacyclohexane-4-spirocyclopentane-2,6-dione (Bocelli, Grenier-Loustalot & Urbanczyk-Lipkowska, 1982). Bonds between C and H atoms (which were refined without constraints) have lengths in the region 0.96–1.09 \AA .

The six-membered ring shows a half-boat conformation; following the Cremer & Pople (1975) notation, the values of the ring-puckering parameters are $Q = 0.630 (5) \text{ \AA}$, $\theta = 46.0 (6)^\circ$ and $\varphi = 2.6 (7)^\circ$. Atoms C(7) and O(1) lie correspondingly 0.89 (1) \AA above and 0.13 (1) \AA below the plane formed by C(1), C(2), C(3) and C(4) atoms. Carbonyl oxygens O(2) and O(3) are above this plane [0.09 (1) and 0.02 (1) \AA respectively]. The striking feature of this conformation is the distinct non-planarity of the anhydride group as a result of the geometric constraints followed from the bicyclic structure of the compound. The bridging atom C(7) is a member of both five- and six-membered rings and the angle C(1)–C(7)–C(4) = $99.2 (2)^\circ$ is far from the standard value indicated for the above-mentioned glutaric anhydrides (about 110°) and is typical for compounds with bicyclo[2.2.1]heptane (Bear & Trotter, 1975; Połński & Dauter, 1986) and bicyclo[3.2.1]octane skeletons (Cheer, Martz, Harpp & Friedlander, 1985). None of the intermolecular distances are shorter than the sum of van der Waals radii. The

additional factor contributing to the non-planarity of the O=C—O—C=O group is probably the repulsion between the methyl at C(7) and the O(1) atom.

The authors thank the Polish Ministry of Science and Higher Education for support from project RP.II.10.

References

- BEAR, C. A. & TROTTER, J. (1975). *Acta Cryst.* B31, 903–904.
 BOCELLI, G. & GRENIER-LOUSTALOT, M. F. (1982). *Acta Cryst.* B38, 2070–2072.
 BOCELLI, G., GRENIER-LOUSTALOT, M. F. & URBANCZYK-LIPKOWSKA, Z. (1982). *J. Crystallogr. Spectrosc. Res.* 12, 407–414.
 CHEER, C. J., MARTZ, E. F., HARPP, D. N. & FRIEDLANDER, B. F. (1985). *Acta Cryst.* C41, 1667–1669.
 CREMER, D. & POPLE, J. A. (1975). *J. Am. Chem. Soc.* 97, 1354–1358.
 HALTZWANGER, R. C., WALBA, D. M. & WAND, M. D. (1980). *Cryst. Struct. Commun.* 9, 1195–1200.
 HAMILTON, W. C. (1959). *Acta Cryst.* 12, 609–610.
 HAŠEK, J. (1985). *Acta Cryst.* C41, 583–586.
 HOEHNE, E. (1966). *Monatsber. Dtsch. Akad. Wiss. Berlin*, 8, 406–410.
International Tables for X-ray Crystallography (1974). Vol. IV. Birmingham: Kynoch Press. (Present distributor D. Reidel, Dordrecht.)
 KOER, F. J., DE HOOG, A. J. & ALTONA, C. (1975). *Recl Trav. Chim. Pays-Bas*, 91, 691–699.
 POŁOŃSKI, T. (1983). *J. Chem. Soc. Perkin Trans. 1*, pp. 305–309.
 POŁOŃSKI, T. & DAUTER, Z. (1986). *J. Chem. Soc., Perkin Trans. 1*, pp. 1781–1788.
 STEWART, J. M. & HALL, S. R. (1983). The XTAL system of crystallographic programs. Tech. Rep. 1364. Univ. of Maryland, College Park, Maryland.

Acta Cryst. (1987). C43, 579–581

Benzyl-9 Diméthyl-5,8 Diméthylamino-3 Carbazolecarbaldéhyde-4*

PAR B. VIOSSAT

Laboratoire de Chimie minérale, UFR de Médecine et de Pharmacie, 34 rue du Jardin des Plantes, 86034 Poitiers CEDEX, France

N. RODIER

Laboratoire de Chimie minérale, Faculté des Sciences Pharmaceutiques et Biologiques, rue J.-B. Clément, 92296 Châtenay-Malabry CEDEX, France

ET C. GANSSER ET C. VIEL

Laboratoire de Chimie thérapeutique, UA 496 CNRS, Faculté des Sciences Pharmaceutiques et Biologiques, rue J.-B. Clément, 92296 Châtenay-Malabry CEDEX, France

(Reçu le 20 avril 1986, accepté le 20 octobre 1986)

Abstract. C₂₄H₂₄N₂O, *M_r* = 356.5, triclinic, *P* $\bar{1}$, *a* = 13.205 (4), *b* = 9.553 (3), *c* = 8.556 (1) Å, α = 104.35 (2), β = 110.11 (2), γ = 98.09 (3)°, *V* = 951.2 Å³, *D_m* = 1.27, *D_x* = 1.24 Mg m⁻³, *Z* = 2, $\lambda(\text{Mo } K\alpha)$ = 0.7107 Å, μ = 0.071 mm⁻¹, *F*(000) = 380, room temperature, *R* = 0.037 for 2663 independent reflections [*I* > 3σ(*I*)]. The carbazole group is not as planar as it is in the carbazole molecule. This distortion is due to steric hindrance. The phenyl ring of the benzyl group and the pyrrole ring are almost perpendicular. The structure can be regarded as consisting either of layers of molecules which are spread out along the (100) planes or of columns built around the [010] axes.

Introduction. La synthèse de la diméthylamino-9 ellipticine par la méthode de Pomeranz (1893) et

Fritsch (1893) conduit à préparer, comme produit intermédiaire, le diméthyl-1,4 diméthylamino-6 carbazole-carbaldéhyde-3. Selon que la formylation, par la méthode de Vilsmeier & Haack (1927), est réalisée sur le diméthyl-1,4 diméthylamino-6 carbazole ou sur son analogue méthoxylé en 6, le groupement CHO se fixe exclusivement en 9 ou de façon prépondérante en 3. La formylation du produit obtenu après blocage de l'azote carbazolique en 9 par le groupement benzyle paraissait donc susceptible de fournir un dérivé formylé en 3. En fait, on obtient un mélange de composés mono et diformylé qui ont été séparés par chromatographie. Par suite de la présence de pics supplémentaires dus aux atomes d'hydrogène aromatiques du groupement benzyle, les spectres de RMN à 250 MHz n'ont pas permis de connaître les positions occupées par le nouveau substituant. C'est la raison pour laquelle l'étude cristallographique a été entreprise. Des cristaux du composé monoformylé ont été obtenus par évaporation

* La numérotation utilisée dans le titre est différente de celle indiquée sur la Fig. 1 et utilisée dans le reste de ce mémoire.